

Citation for published version:

Ji, Y, Bolhuis, A & Watson, ML 2019, 'Staphylococcus aureus products subvert the Burkholderia cenocepacia-induced inflammatory response in airway epithelial cells', *Journal of Medical Microbiology*, vol. 68, no. 12, pp. 1813-1822. <https://doi.org/10.1099/jmm.0.001100>

DOI:

[10.1099/jmm.0.001100](https://doi.org/10.1099/jmm.0.001100)

Publication date:

2019

Document Version

Peer reviewed version

[Link to publication](https://doi.org/10.1099/jmm.0.001100)

© Yuan Ji, Albert Bolhuis, Malcolm L. Watson, 2019. The definitive peer reviewed, edited version of this article is published in *Journal of Medical Microbiology*, Volume 68, issue 12, 2019, <https://doi.org/10.1099/jmm.0.001100>

University of Bath

Alternative formats

If you require this document in an alternative format, please contact:
openaccess@bath.ac.uk

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

***Staphylococcus aureus* products subvert the *Burkholderia cenocepacia*-
induced inflammatory response in airway epithelial cells**

Yuan Ji, Albert Bolhuis, and Malcolm L. Watson*

Department of Pharmacy and Pharmacology, University of Bath, BA2 7AY, United
Kingdom

* Corresponding author's email: prsmw@bath.ac.uk

Keywords: *Staphylococcus aureus*; *Burkholderia cepacia* complex; Host-pathogen
interactions; Immunomodulatory products.

Abstract

Introduction. Polymicrobial infections commonly occur within the lower respiratory tract, although there is a paucity of information regarding the host response to multi-species pathogens. Deleterious effects occur frequently as a result of microbe-microbe interactions, although an unexpected protective effect of *Staphylococcus aureus* has been observed in cystic fibrosis patients co-infected with *Pseudomonas aeruginosa* or species belonging to the *Burkholderia cepacia* complex.

Aim. This study aims to investigate the differences in host responses to mono- and co-infection with *S. aureus* and *B. cenocepacia* in 16HBE human airway epithelial cells, interpret the host receptor and signalling events targeted by *S. aureus*, and characterise staphylococcal immunomodulatory factors.

Methodology and Results. The results show that *B. cenocepacia* activates MAPK and NF- κ B signalling pathways, subsequently eliciting a robust interleukin (IL)-8 production. However, when airway epithelial cells were co-treated with live *B. cenocepacia* bacteria and *S. aureus* supernatants (conditioned medium), the pro-inflammatory response was attenuated. This anti-inflammatory effect was widely exhibited in *S. aureus* isolates tested and was mediated via reduced MAPK and NF- κ B signalling but not via the IL-1 receptor or the tumour necrosis factor receptor modulation. The staphylococcal effectors were characterised as small, heat-stable, non-proteinaceous, and not cell wall-related factors.

Conclusion. This study demonstrates for the first time the host response in a *S. aureus* / *B. cenocepacia* co-infection model and provides insight into a staphylococcal immune evasion mechanism, as well as a therapeutic intervention for excessive inflammation.

Introduction

Lower respiratory tract infections are a major cause of morbidity and mortality globally (Troeger et al., 2017). Individuals with an immune deficiency, chronic obstructive pulmonary disease, or cystic fibrosis (CF) are particularly prone to such respiratory infections (Pragman et al., 2016).

The opportunistic pathogen *Staphylococcus aureus* is one of the major threats that is commonly associated with respiratory infections both in hospital and community environments (Tong et al., 2015). This is not only due to a wide range of virulence factors deployed by *S. aureus* (Haaber et al., 2017; Thammavongsa et al., 2015), but also the rapid emergence of multidrug-resistant strains. *S. aureus* that gains resistance to β -lactam antibiotics as a consequence of adopting the staphylococcal cassette chromosomal *mec* element, becomes methicillin-resistant *S. aureus* (MRSA). This emergence is posing serious challenges to antimicrobial therapy as patients harbouring MRSA are associated with worse clinical prognosis compared to those who never have MRSA (Dasenbrook et al., 2010; Moran et al., 2012).

Burkholderia cepacia complex (Bcc) is a major threat for immunocompromised individuals such as CF patients, as it is transmissible between patients who can develop 'cepacia syndrome' which is a fatal exacerbation of respiratory function as a result of bacteraemia (Hauser and Orsini, 2015; Kenna et al., 2017; LiPuma et al., 1988; Palfreyman et al., 1997). Together with *Pseudomonas aeruginosa* and other species, Bcc are non-fermenting Gram negative bacilli which are not able to utilise glucose in the absence of oxygen. Pathogens of these bacteria have become an increasing challenge in hospital-acquired infections due to the high magnitude of intrinsic antibiotic resistance (Oliveira et al., 2017; Shommu et al., 2015). While the prevalence of *S. aureus* infection is about 60% among CF patients younger than 2

years and peaks between 11 to 17 years old, most Bcc colonisation is acquired after the age of 18 (Cystic Fibrosis Foundation, 2017). Bcc bacteria express a wide range of virulence factors to facilitate the infection process, such as cable pili, flagella, several secretion systems or homologous structures (Leitão et al., 2010; Urban et al., 2005; 2004). In addition, the unusual modification of lipopolysaccharide (LPS) at least partially contributes to the virulence of Bcc. Acylation of lipid A contributes to a much stronger pro-inflammatory response induced by Bcc LPS even compared to LPS from *P. aeruginosa* and *Escherichia coli* (De Soyza et al., 2008). The resulting cytokine storm attracts a large number of immune cells such as neutrophils and mononuclear cells to the lung. If the acute inflammatory response is not resolved, it causes tissue injury which leads to pulmonary fibrosis and possibly develops into a chronic inflammation (Tisoncik et al., 2012).

Chronic pulmonary infection is commonly associated with colonisation with more than one microorganism, but little is known regarding the complicated mechanisms of host-microbe and microbe-microbe interactions and their impact on disease. The microbe-microbe interaction can be synergistic or antagonistic in nature and both forms of interaction can lead to increased antibiotic resistance and virulence (Orazi and O'Toole, 2017; Perez et al., 2014; Ramsey and Whiteley, 2009). Although many studies have reported worse clinical outcomes resulting from microbe-microbe interactions (Bragonzi et al., 2012; Fugère et al., 2014; Korgaonkar et al., 2013), this is not always the case. Contrary to the negative impact by Bcc and *P. aeruginosa*, *S. aureus* co-infection is positively associated with a relatively better lung function, higher survival rate, and delayed lung disease progression (Emerson et al., 2002; Liou et al., 2001; Mayer-Hamblett et al., 2007; Nixon et al., 2001). An *in vivo* study that mimicked the sequential acquisition of early *S. aureus* and late *P. aeruginosa* in the murine lung

shows that pre-infection with *S. aureus* significantly reduced inflammation and the incidence of mortality caused by *P. aeruginosa* but enhanced *P. aeruginosa* persistence (Cigana et al., 2018). We thus hypothesised that the protective effect of *S. aureus* results from dampening the inflammatory response towards Bcc by interfering with the host signalling pathways. Therefore, evaluating whether *S. aureus* colonisation favours the host and understanding the mechanism behind the unexpected protective effect of *S. aureus* may shed light on novel therapies for persistent infections.

In this study, in order to gain insights of the host response to *S. aureus* / *B. cenocepacia* co-colonisation, we assessed the effect of *B. cenocepacia* on the host signalling cascade and inflammatory responses in airway epithelial cells, compared to that of co-infection. Further we identified the host target of staphylococcal immunomodulatory factors, and the nature of immunomodulatory molecules released by *S. aureus*, highlighting the anti-inflammatory modulation of *S. aureus* during Bcc co-infection and potential effects on the clinical outcome.

119 **Material and methods**

120 **Bacteria strains and culture**

121 The bacterial strains used in this study are listed in Table 1. Bacteria were cultured in
122 LB broth (Thermo Fisher Scientific) at 37°C with shaking. Bacterial supernatants were
123 collected and filtered through a 0.2 µm membrane. Where indicated, supernatants
124 were heat-treated by incubating at 95°C for 10 minutes. Heat-killed *S. aureus* were
125 prepared by resuspending the overnight culture to the desired multiplicity of infection
126 (MOI) in DMEM (Gibco, Thermo Fisher Scientific), and heating at 100°C for 1 hour.

127 Proteinase K immobilised on Eupergit® C (Sigma-Aldrich) was used at a concentration
128 of 100 µg/mL in Tris-Cl (50 mM, pH 8) / CaCl₂ (10 mM) at 37°C for 1 hour. At the end
129 of digestion, proteinase K was removed by centrifugation at 250 *g* for 10 minutes.
130 Inactivated proteinase K was prepared by heating at 100°C for 1 hour.

131 Fractionated products of *S. aureus* were prepared by submitting supernatants to 10-
132 kDa- and 3-kDa-cutoff centrifugal concentrators (Vivaspin 20, Sigma-Aldrich),
133 centrifuged at 5000 *g* for 17 minutes and 180 minutes respectively following the
134 manufacturer's instructions. Supernatants were separated into three fractions: >10000
135 molecular weight cut-off (MWCO), 3000-10000 MWCO, and <3000 MWCO. Filtrates
136 and concentrates were adjusted to the original volume with fresh LB, followed by filter
137 sterilisation.

138 **Human airway epithelial cell culture**

139 Human bronchial epithelial 16HBE (obtained from Prof Dieter C. Gruenert, University
140 of California San Francisco) and human airway epithelial Calu-3 (ATCC) were
141 maintained in DMEM, supplemented with 10% heat-inactivated foetal bovine serum,
142 100 units/mL penicillin, and 100 µg/mL streptomycin (all from Gibco, Thermo Fisher

Scientific). Cells were incubated at 37°C in a humidified atmosphere with 5% CO₂. IL-1β and tumour necrosis factor (TNF) α (Peprotech) were used at the concentrations indicated.

Cell viability assay

The cell viability assay with MTT (3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide; Sigma-Aldrich) assay was used as described before (Wachsmann and Lamprecht, 2012). Briefly, 16HBE cells were plated in 96-well plates (5000 cells/well) and incubated for 24 hours. Next, the medium was replaced by 10% *S. aureus* supernatants for 24 hours. After incubating cells with 12 mM MTT for 2 hours, formazan formed in viable cells was dissolved in DMSO for 10 minutes before the optical density was read at 550 nm.

Detection of Cytokine Production

At the end of stimulation, conditioned medium of 16HBE cells was collected and the concentration of interleukin (IL)-8 was quantified using Maxisorp ELISA plates (Nunc, Thermo Fisher Scientific) and a commercially available ELISA kit (eBioscience, Thermo Fisher Scientific) according to the manufacturer's instructions.

Western Blot Analysis

Following stimulation, cell lysates were prepared and the concentration of total protein was quantified using PierceTM BCA Protein Assay Kit (Thermo Fisher Scientific). Protein samples were resolved on an SDS-PAGE gel, and immunoblotted on a nitrocellulose membrane (Bio-Rad). The membrane was probed with antibodies anti phospho-NF-κB-p65, phospho-p38, phospho-Erk, and p38 (Cell Signaling Technology), followed by incubation with secondary antibodies (Li-Cor Biosciences).

Quantitative analysis of the signal was performed using the Li-Cor Odyssey Clx imaging system.

Statistical Analysis

Data were expressed as the mean \pm standard error of the mean (SEM) for $n=3$ independent biological repeats. Results were compared using Graphpad Prism and one-way or two-way ANOVA, followed by Tukey or Dunnett's multiple comparisons test, with $p<0.05$ considered significant.

Results

***B. cenocepacia*-stimulated IL-8 response is suppressed by *S. aureus* products.**

IL-8 plays an important role at the early stage of infection by chemoattracting granulocytes such as neutrophils (Perret et al., 2012). Hence, the level of IL-8 is a biomarker of pro-inflammatory responses and inflammation (Dennehy et al., 2017; Palfreyman et al., 1997). *B. cenocepacia* induced a robust increase in IL-8 response in 16HBE cells (Figure 1A). Although supernatants (in LB medium) of *S. aureus* MRSA252 or MSSA NCTC 6571 did not change IL-8 production compared to the control, they significantly downregulated IL-8 induced by *B. cenocepacia* (23% and 31% less than the *B. cenocepacia*-treated group, respectively). In contrast, heat-killed *S. aureus* did not have the suppressing effect (Figure 1B).

The same experiments were done with a collection of *S. aureus* isolates, either supernatants (Figure 1C). Supernatants from all the tested isolates downregulated the level of IL-8 production by different amounts (5.4%-38.0%). Although some of the differences were small, consistent changes were seen within each experiment and these were statistically significant ($p<0.05$). The ability of heat-killed bacteria to inhibit *B. cenocepacia*-elicited IL-8 production (Figure 1D). However, whole heat-killed bacteria had no effect on *B. cenocepacia*-induced IL-8 production. These data indicate that anti-inflammatory activity in these experiments is expressed in secreted products but not heat-stable *S. aureus* cell surface structures. This contrasts with the nasal isolate *S. aureus* anti-inflammatory activity reported by Peres et al. (2015), which was exhibited by whole bacteria subjected to the same heat treatment conditions.

To confirm the suppressing effect is not due to cell damage, an MTT assay was used to assess cell viability (Figure 1E). For the majority of *S. aureus* strains, their

supernatants did not cause epithelial cell damage, indicating that there was no causality between IL-8 suppression and induction of epithelial cell damage. Exceptions were SH1000 and RN4220 which did reduce cell viability and, interestingly, these alone caused a doubling in IL-8 production (Figure 1C).

MSSA NCTC 6571 products block MAPK p38 and Erk, and NF- κ B signalling activated by *B. cenocepacia*

The recognition of conserved microbial structures of pathogens by pattern recognition receptors leads to activation of MAPK and NF- κ B signalling pathways, subsequently initiating the expression of pro-inflammatory genes to mount an immune response (Pandey et al., 2014). NF- κ B and MAPK pathways contribute to IL-8 expression at multiple levels (Hoffmann et al., 2002). Since *B. cenocepacia* induced a strong IL-8 production, the effects of *B. cenocepacia* on MAPK and NF- κ B signalling in 16HBE cells were assessed. Cells were challenged with *B. cenocepacia* at an MOI of 5 from 5 minutes to 180 minutes (Figure 2 A-D). For the MAPK family, phosphorylation of p38 was maximal at 120 minutes (200% more than time 0), while phosphorylation of Erk peaked at 5 minutes (156% more than time 0), followed by a decrease and then a secondary rise at 120 minutes (133% more than time 0). For NF- κ B signalling, *B. cenocepacia* upregulated phosphorylation of p65, particularly after 120 minutes.

In order to study the modulation on host signalling by this co-stimulation model, 16HBE cells were treated with *B. cenocepacia* (5 MOI) and 10% *S. aureus* supernatant for 2 hours (Figure 2E-H). *B. cenocepacia* led to phosphorylation of p38, Erk, and p65, whereas *S. aureus* supernatant alone did not change any tested signalling events. Notably, MSSA NCTC 6571 but not MRSA252 supernatant reduced phosphorylation of p38, Erk and p65 stimulated by *B. cenocepacia*. The reduction in p38 and p65 signalling was significant compared with *B. cenocepacia* alone ($p<0.05$). This strain

selectivity is consistent with the differential inhibition of the pro-inflammatory response (Figure 1A) by MSSA NTCT 6571 and MRSA252. Therefore, these findings suggest that MSSA NCTC 6571 supernatant may suppress the pro-inflammatory effect of *B. cenocepacia* by inhibiting MAPK and NF- κ B signalling.

Characterisation of anti-inflammatory factors in *S. aureus* supernatant

To investigate the nature of anti-inflammatory components, *S. aureus* supernatants were heated at 95°C for 10 minutes. These heat-treated supernatants were still able to inhibit the *B. cenocepacia*-stimulated IL-8 response (Figure 3A), indicating that the components are heat stable.

Further, *S. aureus* supernatants were treated with solid-phase proteinase K for 1 hour at 37°C before removal to avoid any effect of the proteinase on released IL-8. Proteinase K-digested supernatants did not restore IL-8 secretion caused by *B. cenocepacia*; rather, IL-8 secretion was nearly reduced to the baseline (Figure 3B). Further, the 16HBE cultures incubated with proteinase-treated supernatant alone did not inhibit basal IL-8 levels (Figure 3B), indicating that no amounts of protease K was carried over into the production assay. Heat-inactivated proteinase K-treated supernatants displayed similar inhibition level as untreated supernatants (Figure 1A), indicating the components are either non-proteinaceous or proteinase K-resistant.

To assess the approximate molecular weight of the secretory *S. aureus* components causing a reduction in *B. cenocepacia*-induced IL-8 production, *S. aureus* supernatants were separated into different molecular cut-off fractions using selective membranes (Figure 3C). Components in the supernatants with a molecular weight less than 3 kDa, but not above 3 kDa reduced IL-8 production back to the same level as the control, which indicates the anti-inflammatory factors are small molecules.

As staphylococcal virulence factors (such as SpA and Panton-Valentine Leucocidin) target the IL-1 receptor (IL-1R) and tumour necrosis factor receptor (TNFR), leading to IL-8 expression (Gomez et al., 2004; Labrousse et al., 2014), in order to elucidate whether the *S. aureus*-induced anti-inflammatory response was selective for bacterial stimuli, 16HBE cells were treated with IL-1 β and TNF α , resulting in the same amount of IL-8 as the *B. cenocepacia*-treated group (Figure 3D). However, contrary to the reduction seen with *S. aureus* supernatant and J2315 co-stimulation, *S. aureus* supernatants did not suppress IL-1 β / TNF α -induced IL-8 production, indicating that IL-1R or TNFR signalling is not associated with this process and the suppressive effect appears to be targeted toward pathways involved in the *B. cenocepacia* IL-8 response.

Discussion

This study is the first to demonstrate the host response caused by *S. aureus* / *B. cenocepacia* co-infection in airway epithelial cells. We demonstrate that *S. aureus* secreted products, but not cell wall-related components are able to suppress the pro-inflammatory response induced by *B. cenocepacia*. The active compound(s) are primarily characterised as small, heat-stable, non-proteinaceous, and targeting MAPK and NF- κ B signalling.

Previous studies have concentrated on the interactions between *S. aureus* and *P. aeruginosa* as they are two prevalent pathogens involved in CF lung infections (Ahlgren et al., 2015; Baldan et al., 2014; Chekabab et al., 2015; DeLeon et al., 2014; Fugère et al., 2014; Orazi and O'Toole, 2017). However, the impact of *S. aureus* / *Bcc* interactions on host responses has never been assessed. Herein, we demonstrate that in the context of *S. aureus* / *B. cenocepacia* co-infection, supernatants from *S. aureus* isolates subverted *B. cenocepacia* live bacteria-induced IL-8 production in airway epithelial cells, suggesting that *S. aureus* releases anti-inflammatory compounds into the extracellular milieu. This effect of *S. aureus* provides a possible explanation of the protective effect of *S. aureus* colonisation on lung function and patient survival (Liou et al., 2001; Mayer-Hamblett et al., 2007).

Interestingly, all the supernatants from *S. aureus* isolates tested were able to subvert *B. cenocepacia*-induced IL-8 production ($p < 0.05$). Even hyperinflammatory isolates such as SH1000 and RN4220, where the supernatants induced strong IL-8 production (Figure 1C), were able to reduce the inflammatory response, indicating the components may be widely expressed in *S. aureus*. The secretome of *S. aureus* involves a large arsenal of immune evasion factors that inhibit complement, chemotaxis, phagocytosis, and induce neutrophil lysis. A whole genome study divided

308 88 *S. aureus* isolates into 25 clonal complex lineages and investigated 43 immune
309 evasion genes (McCarthy and Lindsay, 2013). The results revealed that although
310 genes encoding leucocidin are highly conserved, there are variants among other
311 genes, indicating each lineage applies different strategies and functions of immune
312 evasion.

313 The MAPK family and NF- κ B signalling pathways are known to play key roles in the
314 host response to bacterial pathogens by regulating the expression of cytokines and
315 chemokines (Krachler et al., 2011). However, pathogens have evolved to exploit these
316 pathways. For example, *B. cenocepacia* K56-2 can significantly upregulate
317 phosphorylation of Akt, Erk, and NF- κ B in 16HBE cells within 30 minutes of stimulation
318 (Gillette et al., 2013). In parallel, *B. cenocepacia* BC7 activates phosphorylation of p38
319 and JNK in CF bronchial epithelial IB3 cells, whereas these pathways remain
320 unphosphorylated in contact with an environmental *B. cepacia* strain (Sajjan et al.,
321 2008). As K56-2 and BC7 both belong to the highly virulent ET-12 lineage, this
322 indicates that hyperactivation of host signalling is related to more severe lung
323 inflammation caused by *B. cenocepacia* isolates.

324 In our model, *B. cenocepacia* caused phosphorylation of p38, Erk, and p65 at 120
325 minutes post-infection. Of note, after a transient increase at 5 minutes post-infection,
326 phosphorylation of Erk peaked again at 120 minutes. The similar bi-phasic activation
327 pattern of Erk and p38 was observed in the response to *B. pseudomallei* both *in vitro*
328 and *in vivo* (D'Elia et al., 2017). Thirty-minutes post-infection, murine alveolar
329 macrophages exhibited transient phosphorylation of p38 and Erk1, and at 2 h post-
330 infection exhibited sustained activation. As Erk regulates apoptosis in response to
331 oxidative stress (Li et al., 2016), D'Elia et al. (2017) proposed that the transient
332 activation at the early stage is associated with host defence, while during the latter

333 stage of infection host cells might initiate apoptosis as indicated by continuous Erk
334 activation, and thereby *B. pseudomallei* gains time for intracellular proliferation by
335 delaying apoptosis through modulate the dynamics of Erk activation.

336 Upon co-stimulation with *B. cenocepacia* live bacteria and *S. aureus* supernatant,
337 activation of MAPK and NF- κ B was assessed where MSSA NCTC 6571 supernatant
338 abolished phosphorylation of p38, Erk, and p65 caused by *B. cenocepacia*. MRSA252
339 supernatant did not affect either baseline or *B. cenocepacia*-induced phosphorylation,
340 which correlates with its lower inhibitory ability on IL-8 production compared to MSSA
341 NCTC 6571. Our results are in partial agreement with the results displayed by
342 Chekabab et al., (2015) as they showed *S. aureus* supernatants inhibited *P.*
343 *aeruginosa* supernatant-induced inflammatory response through blocking NF- κ B
344 activation but not p38. However, Chekabab et al. (2015) stimulated airway epithelial
345 Beas-2B cells with synthetic Toll-like receptor (TLR)-2/1 agonist Pam₃CSK4 but not *P.*
346 *aeruginosa*, but bacteria themselves contain far more complicated ligands which could
347 potentially activate the MAPK family. Additionally, the downstream signalling cascades
348 engaged in the anti-inflammatory response may be cell type-specific. For example, in
349 human peripheral blood mononuclear cells, among a group of nasal *S. aureus* isolates,
350 some isolates induce anti-inflammatory IL-10 production through PI3K/Akt/mTOR and
351 Erk signalling whereas some induce pro-inflammatory TNF α through a p38-dependent
352 pathway (Peres et al., 2015). This suggests that *S. aureus* manipulates different
353 signalling cascades to initiate immune tolerance through heterogenicity expression of
354 ligands in the process of infection.

355 In further experiments, the treatments of heat, proteinase K, and different molecular
356 weight cut-off fractions were examined in order to determine the nature of the anti-
357 inflammatory compounds in *S. aureus* supernatants. Consistent with previous studies,

the compounds were characterised as small (<3 kDa) and heat-resistant (Chekabab et al., 2015; Tajima et al., 2006). As secondary or tertiary structures of proteins are usually disrupted during boiling, the staphylococcal factors are unlikely to be proteins. Similarly, when co-stimulated with *B. cenocepacia*, IL-8 production elicited by proteinase K-treated *S. aureus* supernatants showed a 50% decrease compared to deactivated proteinase K-treated *S. aureus* supernatants, indicating that anti-inflammatory components are less likely to be peptides due to the nonspecific protease activity of proteinase K. A possible explanation is that proteinaceous pro-inflammatory components in *S. aureus* supernatants were degraded by proteinase K and thus the anti-inflammatory effectors overpowered the pro-inflammatory counterparts. For example, the major virulence factor SpA not only contributes to immune evasion by binding to immunoglobulin, but also elicits a strong pro-inflammatory response through interacting with tumour necrosis factor receptor 1 and epidermal growth factor receptor (Gomez et al., 2004; Graille et al., 2000; Soong et al., 2011). So far only one research group reported small, heat-, and protease-resistant molecules derived from MRSA and *S. epidermidis* conditioned medium that induce nitric oxide in human upper airway epithelial cells (Carey et al., 2016; 2015). Although these studies did not assess production of cytokines and chemokines in response to the factors, the result strongly indicates the existence of small, heat- and protease-resistant molecules that are secreted by *S. aureus*.

We also investigated whether *S. aureus* factors interact with other receptors such as IL-1R and TNFR that promote the inflammatory response. The results showed that the inhibitory effect of *S. aureus* secreted products was *B. cenocepacia* stimuli-specific, as *S. aureus* supernatants had no effect on IL-1 β / TNF α -induced IL-8 production. This is not surprising as *S. aureus* can colonise human airway asymptotically, and

therefore is able to trigger a low level of inflammatory response and even suppress it in order to induce immune tolerance. Known staphylococcal factors that antagonise the vital microbial sensor TLRs, include staphylococcal superantigen-like protein 3 and TIR domain protein targeting TLR2 (Askarian et al., 2014; Koymans et al., 2017), and phenol-soluble modulin α 1-3 targeting TLR4 (Chu et al., 2018). However, little is known about whether the immunosuppression of *S. aureus* is beneficial to the host. To further explore our findings, the impact of *S. aureus* on *B. cenocepacia* infection need to be investigated using an *in vivo* respiratory infection model. Our findings for the first time illustrate the host response to *S. aureus* and *B. cenocepacia* co-infection in the airway epithelial cells, and reveal staphylococcal anti-inflammatory factors that suppress *B. cenocepacia*-induced IL-8 production through blocking MAPK and NF- κ B signalling. To conclude, we reveal immune evasion factors secreted by *S. aureus* that are small, heat-stable, and non-proteinaceous, which reduce the pro-inflammatory response induced by CF-isolated *B. cenocepacia* in 16HBE cells. These results suggest a potential treatment for patients with chronic pulmonary infections, whereby drugs targeting MAPK and NF- κ B pathways could be applied to reduce inflammation without facilitating infections.

Author statement

The authors declare that there are no conflicts of interest.

Acknowledgement

YJ was funded by a University of Bath Scholarship.

407 **References**

- 408 Ahlgren, H.G., Benedetti, A., Landry, J.S., Bernier, J., Matouk, E., Radzioch, D., Lands, L.C.,
409 Rousseau, S., Nguyen, D., 2015. Clinical outcomes associated with *Staphylococcus aureus*
410 and *Pseudomonas aeruginosa* airway infections in adult cystic fibrosis patients. *BMC Pulm*
411 *Med* 15, 918. doi: org/10.1186/s12890-015-0062-7.
- 412 Askarian, F., van Sorge, N.M., Sangvik, M., Beasley, F.C., Henriksen, J.R., Sollid, J.U.E.,
413 van Strijp, J.A.G., Nizet, V., Johannessen, M., 2014. A *Staphylococcus aureus* TIR domain
414 protein virulence factor blocks TLR2-mediated NF- κ B signaling. *J Innate Immun* 6, 485–498.
415 doi: 10.1159/000357618.
- 416 Baba, T., Bae, T., Schneewind, O., Takeuchi, F., Hiramatsu, K., 2008. Genome sequence of
417 *Staphylococcus aureus* strain Newman and comparative analysis of staphylococcal
418 genomes: Polymorphism and evolution of two major pathogenicity islands. *J Bacterio* 190,
419 300–310. doi: 10.1128/JB.01000-07.
- 420 Baldan, R., Cigana, C., Testa, F., Bianconi, I., De Simone, M., Pellin, D., Di Serio, C.,
421 Bragonzi, A., Cirillo, D.M., 2014. Adaptation of *Pseudomonas aeruginosa* in Cystic Fibrosis
422 airways influences virulence of *Staphylococcus aureus* *in vitro* and murine models of co-
423 infection. *PLoS ONE* 9, e89614. doi: 10.1371/journal.pone.0089614.
- 424 Bragonzi, A., Farulla, I., Paroni, M., Twomey, K.B., Pirone, L., Lorè, N.I., Bianconi, I.,
425 Dalmastri, C., Ryan, R.P., Bevivino, A., 2012. Modelling co-infection of the cystic fibrosis
426 lung by *Pseudomonas aeruginosa* and *Burkholderia cenocepacia* reveals influences on
427 biofilm formation and host response. *PLoS ONE* 7, e52330. doi:
428 10.1371/journal.pone.0052330.
- 429 Carey, R.M., Chen, B., Adappa, N.D., Palmer, J.N., Kennedy, D.W., Lee, R.J., Cohen, N.A.,
430 2016. Human upper airway epithelium produces nitric oxide in response to *Staphylococcus*
431 *epidermidis*. *Int Forum Allergy Rhinol* 6, 1238–1244. doi: 10.1002/alr.21837.
- 432 Carey, R.M., Workman, A.D., Chen, B., Adappa, N.D., Palmer, J.N., Kennedy, D.W., Lee,
433 R.J., Cohen, N.A., 2015. *Staphylococcus aureus* triggers nitric oxide production in human
434 upper airway epithelium. *Int Forum Allergy Rhinol* 5, 808–813. doi: 10.1002/alr.21568.
- 435 Chekabab, S.M., Silverman, R.J., Lafayette, S.L., Luo, Y., Rousseau, S., Nguyen, D., 2015.
436 *Staphylococcus aureus* inhibits IL-8 responses induced by *Pseudomonas aeruginosa* in
437 airway epithelial cells. *PLoS ONE* 10, e0137753–19. doi: org/10.1371/journal.pone.0137753.
- 438 Chu, M., Zhou, M., Jiang, C., Chen, X., Guo, L., Zhang, M., Chu, Z., Wang, Y., 2018.
439 *Staphylococcus aureus* phenol-soluble modulins α 1– α 3 act as novel Toll-like receptor (TLR)
440 4 antagonists to inhibit HMGB1/TLR4/NF- κ B signaling pathway. *Front Immunol* 9, 596–13.
441 doi: 10.3389/fimmu.2018.00862.
- 442 Cigana, C., Bianconi, I., Baldan, R., De Simone, M., Riva, C., Sipione, B., Rossi, G., Cirillo,
443 D.M., Bragonzi, A., 2018. *Staphylococcus aureus* impacts *Pseudomonas aeruginosa* chronic
444 respiratory disease in murine models. *J Infect Dis* 217, 933–942. doi: 10.1093/infdis/jix621.
- 445 Collins, J., Rudkin, J., Recker, M., Pozzi, C., Gara, J.P.O.A., Massey, R.C., 2010. Offsetting
446 virulence and antibiotic resistance costs by MRSA. *ISME J* 4, 577–584. doi:
447 10.1038/ismej.2009.151.

448 Cystic Fibrosis Foundation, 2017. Cystic Fibrosis Foundation Patient Registry 2016 Annual
449 Data Report. Bethesda, Maryland.

450 Dasenbrook, E.C., Checkley, W., Merlo, C.A., Konstan, M.W., Lechtzin, N., Boyle, M.P.,
451 2010. Association between respiratory tract methicillin-resistant *Staphylococcus aureus* and
452 survival in cystic fibrosis. *JAMA* 303, 2386–2392. doi: 10.1001/jama.2010.791.

453 De Soyza, A., Silipo, A., Lanzetta, R., Govan, J.R., Molinaro, A., 2008. Chemical and
454 biological features of *Burkholderia cepacia* complex lipopolysaccharides. *Innate Immun* 14,
455 127–144. doi: 10.1177/1753425908093984.

456 DeLeon, S., Clinton, A., Fowler, H., Everett, J., Horswill, A.R., Rumbaugh, K.P., 2014.
457 Synergistic interactions of *Pseudomonas aeruginosa* and *Staphylococcus aureus* in an *in*
458 *vitro* wound model. *Infect Immun* 82, 4718–4728. doi: 10.1128/IAI.02198-14.

459 Dennehy, R., Romano, M., Ruggiero, A., Mohamed, Y.F., Dignam, S.L., Mujica Troncoso,
460 C., Callaghan, M., Valvano, M.A., Berisio, R., McClean, S., 2017. The *Burkholderia*
461 *cenocepacia* peptidoglycan-associated lipoprotein is involved in epithelial cell attachment
462 and elicitation of inflammation. *Cell Microbiol* 19(5): e12691. doi: 10.1111/cmi.12691.

463 D'Elia, R.V., Saint, R.J., Newstead, S.L., Clark, G.C., Atkins, H.S., 2017. Mitogen-activated
464 protein kinases (MAPKs) are modulated during *in vitro* and *in vivo* infection with the
465 intracellular bacterium *Burkholderia pseudomallei*. *Eur J Clin Microbiol Infect Dis* 36, 2147–
466 2154. doi: 10.1007/s10096-017-3038-0.

467 Emerson, J., Rosenfeld, M., McNamara, S., Ramsey, B., Gibson, R.L., 2002. *Pseudomonas*
468 *aeruginosa* and other predictors of mortality and morbidity in young children with cystic
469 fibrosis. *Pediatr Pulmonol* 34, 91–100. doi: 10.1002/ppul.10127

470• Fugère, A., Lalonde Séguin, D., Mitchell, G., Déziel, E., Dekimpe, V., Cantin, A.M., Frost, E.,
471 Malouin, F., 2014. Interspecific small molecule interactions between clinical isolates of
472 *Pseudomonas aeruginosa* and *Staphylococcus aureus* from adult cystic fibrosis patients.
473 *PLoS ONE* 9, e86705. doi.org/10.1371/journal.pone.0086705

474 Gillette, D.D., Shah, P.A., Cremer, T., Gavrilin, M.A., Besecker, B.Y., Sarkar, A., Knoell,
475 D.L., Cormet-Boyaka, E., Wewers, M.D., Butchar, J.P., Tridandapani, S., 2013. Analysis of
476 human bronchial epithelial cell proinflammatory response to *Burkholderia cenocepacia*
477 infection. *J Biol Chem* 288, 3691–3695. doi: 10.1074/jbc.C112.430298.

478 Gomez, M.I., Lee, A., Reddy, B., Muir, A., Soong, G., Pitt, A., Cheung, A., Prince, A., 2004.
479 *Staphylococcus aureus* protein A induces airway epithelial inflammatory responses by
480 activating TNFR1. *Nat Med* 10, 842–848. doi: 10.1038/nm1079.

481 Graille, M., Stura, E.A., Corper, A.L., Sutton, B.J., Taussig, M.J., Charbonnier, J.B.,
482 Silverman, G.J., 2000. Crystal structure of a *Staphylococcus aureus* protein A domain
483 complexed with the Fab fragment of a human IgM antibody: structural basis for recognition
484 of B-cell receptors and superantigen activity. *PNAS* 97, 5399–5404. doi:
485 10.1073/pnas.97.10.5399.

486 Haaber, J., Penadés, J.R., Ingmer, H., 2017. Transfer of antibiotic resistance in
487 *Staphylococcus aureus*. *Trends Microbiol* 25, 893–905. doi: 10.1016/j.tim.2017.05.011.

488 Hauser, N., Orsini, J., 2015. Cepacia syndrome in a non-cystic fibrosis patient. *Case Rep*
489 *Infect Dis* 2015, 537627–4. doi: 10.1155/2015/537627.

490 Hoffmann, E., Dittrich-Breiholz, O., Holtmann, H., Kracht, M., 2002. Multiple control of
491 interleukin-8 gene expression. *J Leukoc Biol* 72, 847–855. doi.org/10.1189/jlb.72.5.847.

492 Holden, M.T.G., Feil, E.J., Lindsay, J.A., Peacock, S.J., Day, N.P.J., Enright, M.C., Foster,
493 T.J., Moore, C.E., Hurst, L., Atkin, R., Barron, A., Bason, N., Bentley, S.D., Chillingworth, C.,
494 Chillingworth, T., Churcher, C., Clark, L., Corton, C., Cronin, A., Doggett, J., Dowd, L.,
495 Feltwell, T., Hance, Z., Harris, B., Hauser, H., Holroyd, S., Jagels, K., James, K.D., Lennard,
496 N., Line, A., Mayes, R., Moule, S., Mungall, K., Ormond, D., Quail, M.A., Rabbino-witsch, E.,
497 Rutherford, K., Sanders, M., Sharp, S., Simmonds, M., Stevens, K., Whitehead, S., Barrell,
498 B.G., Spratt, B.G., Parkhill, J., 2004. Complete genomes of two clinical *Staphylococcus*
499 *aureus* strains: evidence for the rapid evolution of virulence and drug resistance. *PNAS* 101,
500 9786–9791. doi: 10.1073/pnas.0402521101.

501 Holden, M.T.G., Seth-Smith, H.M.B., Crossman, L.C., Sebaihia, M., Bentley, S.D., Cerdeno-
502 Tarraga, A.M., Thomson, N.R., Bason, N., Quail, M.A., Sharp, S., Cherevach, I., Churcher,
503 C., Goodhead, I., Hauser, H., Holroyd, N., Mungall, K., Scott, P., Walker, D., White, B.,
504 Rose, H., Iversen, P., Mil-Homens, D., Rocha, E.P.C., Fialho, A.M., Baldwin, A., Dowson,
505 C., Barrell, B.G., Govan, J.R., Vandamme, P., Hart, C.A., Mahenthiralingam, E., Parkhill, J.,
506 2009. The Genome of *Burkholderia cenocepacia* J2315, an epidemic pathogen of cystic
507 fibrosis patients. *J Bacterio* 191, 261–277. doi: 10.1128/JB.01230-08.

508 Kenna, D.T.D., Lilley, D., Coward, A., Martin, K., Perry, C., Pike, R., Hill, R., Turton, J.F.,
509 2017. Prevalence of *Burkholderia* species, including members of *Burkholderia cepacia*
510 complex, among UK cystic and non-cystic fibrosis patients. *J Med Microbiol* 66, 490–501.
511 doi: 10.1099/jmm.0.000458.

512 Korgaonkar, A., Trivedi, U., Rumbaugh, K.P., Whiteley, M., 2013. Community surveillance
513 enhances *Pseudomonas aeruginosa* virulence during polymicrobial infection. *PNAS* 110,
514 1059–1064. doi: 10.1073/pnas.1214550110.

515 Koymans, K.J., Goldmann, O., Karlsson, C.A.Q., Sital, W., Thänert, R., Bisschop, A.,
516 Vrieling, M., Malmström, J., van Kessel, K.P.M., de Haas, C.J.C., van Strijp, J.A.G., Medina,
517 E., 2017. The TLR2 antagonist staphylococcal superantigen-like protein 3 acts as a
518 virulence factor to promote bacterial pathogenicity *in vivo*. *J Innate Immun* 9, 561–573. doi:
519 10.1159/000479100.

520 Krachler, A.M., Woolery, A.R., Orth, K., 2011. Manipulation of kinase signaling by bacterial
521 pathogens. *J Cell Biol* 195, 1083–1092. doi: 10.1083/jcb.201107132.

522 Labrousse, D., Perret, M., Hayez, D., Da Silva, S., Badiou, C., Couzon, F., Bès, M.,
523 Chavanet, P., Lina, G., Vandenesch, F., Croisier-Bertin, D., Henry, T., 2014. Kineret®/IL-1ra
524 blocks the IL-1/IL-8 inflammatory cascade during recombinant Panton Valentine leukocidin-
525 triggered pneumonia but not during *S. aureus* infection. *PLoS ONE* 9, e97546. doi:
526 10.1371/journal.pone.0097546.

527 Leitão, J.H., Sousa, S.A., Ferreira, A.S., Ramos, C.G., Silva, I.N., Moreira, L.M., 2010.
528 Pathogenicity, virulence factors, and strategies to fight against *Burkholderia cepacia*
529 complex pathogens and related species. *Appl Microbiol Biotechnol* 87, 31–40. doi:
530 10.1007/s00253-010-2528-0.

531 Li, Y., Jiang, Z., Xue, D., Deng, G., Li, M., Liu, X., Wang, Y., 2016. *Mycoplasma*
532 *ovipneumoniae* induces sheep airway epithelial cell apoptosis through an ERK signalling-
533 mediated mitochondria pathway. *BMC Microbiol* 16, 222. doi: 10.1186/s12866-016-0842-0.

534 Liou, T.G., Adler, F.R., Fitzsimmons, S.C., Cahill, B.C., Hibbs, J.R., Marshall, B.C., 2001.
535 Predictive 5-year survivorship model of cystic fibrosis. *Am J Epidemiol* 153, 345–352. doi:
536 10.1093/aje/153.4.345.

537 LiPuma, J.J., Mortensen, J.E., Dasen, S.E., Edlind, T.D., Schidlow, D.V., Burns, J.L., Stull,
538 T.L., 1988. Ribotype analysis of *Pseudomonas cepacia* from cystic fibrosis treatment
539 centers. *J Pediatr* 113, 859–862. doi: 10.1016/s0022-3476(88)80018-0.

540 Mayer-Hamblett, N., Aitken, M.L., Accurso, F.J., Kronmal, R.A., Konstan, M.W., Burns, J.L.,
541 Sagel, S.D., Ramsey, B.W., 2007. Association between pulmonary function and sputum
542 biomarkers in cystic fibrosis. *Am J Respir Crit Care Med* 175, 822–828. doi:
543 10.1164/rccm.200609-1354OC.

544 McCarthy, A.J., Lindsay, J.A., 2013. *Staphylococcus aureus* innate immune evasion is
545 lineage-specific: a bioinformatics study. *Infect Genet Evol* 19, 7–14. doi:
546 10.1016/j.meegid.2013.06.012.

547 Moran, G.J., Krishnadasan, A., Gorwitz, R.J., Fosheim, G.E., Albrecht, V., Limbago, B.,
548 Talan, D.A., 2012. Prevalence of methicillin-resistant *Staphylococcus aureus* as an etiology
549 of community-acquired pneumonia. *Clin Infect Dis* 54, 1126–1133. doi: 10.1093/cid/cis022.

550 Nair, D., Memmi, G., Hernandez, D., Bard, J., Beaume, M., Gill, S., Francois, P., Cheung,
551 A.L., 2011. Whole-genome sequencing of *Staphylococcus aureus* strain RN4220, a key
552 laboratory strain used in virulence research, identifies mutations that affect not only virulence
553 factors but also the fitness of the strain. *J Bacterio* 193, 2332–2335. doi: 10.1128/JB.00027-
554 11.

555 Nixon, G.M., Armstrong, D.S., Carzino, R., Carlin, J.B., Olinsky, A., Robertson, C.F.,
556 Grimwood, K., 2001. Clinical outcome after early *Pseudomonas aeruginosa* infection in
557 cystic fibrosis. *J Pediatr* 138, 699–704. doi: 10.1067/mpd.2001.112897.

558 Oliveira, M.E.F., Araújo, D.G., Oliveira, S.R., 2017. Resistance of non-fermenting Gram-
559 negative bacilli isolated from blood cultures from an emergency hospital. *J Bras Patol Med*
560 *Lab* 53, 87–91. doi.org/10.5935/1676-2444.20170013.

561 Orazi, G., O'Toole, G.A., 2017. *Pseudomonas aeruginosa* alters *Staphylococcus aureus*
562 sensitivity to vancomycin in a biofilm model of cystic fibrosis infection. *MBio* 8, 4 e00873-17.
563 doi.org/10.1128/mBio.00873-17.

564 O'Neill, A.J., 2010. *Staphylococcus aureus* SH1000 and 8325-4: comparative genome
565 sequences of key laboratory strains in staphylococcal research. *Lett Appl Microbiol* 51, 358–
566 361. doi: 10.1111/j.1472-765X.2010.02885.x.

567 Palfreyman, R.W., Watson, M.L., Eden, C., Smith, A.W., 1997. Induction of Biologically
568 Active Interleukin-8 from lung epithelial cells by *Burkholderia (Pseudomonas) cepacia*
569 products. *Infect Immun* 65, 617–622.

570 Pandey, S., Kawai, T., Akira, S., 2014. Microbial sensing by Toll-like receptors and
571 intracellular nucleic acid sensors. *Cold Spring Harb Perspect Biol* 7, a016246–a016246. doi:
572 10.1101/cshperspect.a016246.

573 Perret, M., Badiou, C., Lina, G., Burbaud, S., Benito, Y., Bès, M., Cottin, V., Couzon, F.,
574 Juruj, C., Dauwalder, O., Goutagny, N., Diep, B.A., Vandenesch, F., Henry, T., 2012. Cross-
575 talk between *Staphylococcus aureus* leukocidins-intoxicated macrophages and lung
576 epithelial cells triggers chemokine secretion in an inflammasome-dependent manner. *Cell*
577 *Microbiol* 14, 1019–1036. doi: 10.1111/j.1462-5822.2012.01772.x.

578 Peres, A.G., Stegen, C., Li, J., Xu, A.Q., Levast, B., Surette, M.G., Cousineau, B.,
579 Desrosiers, M., Madrenas, J., 2015. Uncoupling of pro- and anti-inflammatory properties of
580 *Staphylococcus aureus*. *Infect Immun* 83, 1587–1597. doi: 10.1128/IAI.02832-14.

581 Perez, A.C., Pang, B., King, L.B., Tan, L., Murrah, K.A., Reimche, J.L., Wren, J.T.,
582 Richardson, S.H., Ghandi, U., Swords, W.E., 2014. Residence of *Streptococcus*
583 *pneumoniae* and *Moraxella catarrhalis* within polymicrobial biofilm promotes antibiotic
584 resistance and bacterial persistence *in vivo*. *Pathog Dis* 70, 280–288. doi: 10.1111/2049-
585 632X.12129.

586 Pragman, A.A., Berger, J.P., Williams, B.J., 2016. Understanding persistent bacterial lung
587 infections: clinical implications informed by the biology of the microbiota and biofilms. *Clin*
588 *Pulm Med* 23, 57–66. doi: 10.1097/cpm.000000000000108.

589 Ramsey, M.M., Whiteley, M., 2009. Polymicrobial interactions stimulate resistance to host
590 innate immunity through metabolite perception. *PNAS* 106, 1578–1583. doi:
591 10.1073/pnas.0809533106.

592 Sajjan, U.S., Hershenson, M.B., Forstner, J.F., LiPuma, J.J., 2008. *Burkholderia*
593 *cenoecepacia* ET12 strain activates TNFR1 signalling in cystic fibrosis airway epithelial cells.
594 *Cell Microbiol* 10, 188–201. doi: 10.1111/j.1462-5822.2007.01029.x.

595 Shommu, N.S., Vogel, H.J., Storey, D.G., 2015. Potential of metabolomics to reveal
596 *Burkholderia cepacia* complex pathogenesis and antibiotic resistance. *Front Microbio* 6, 668.
597 doi: 10.3389/fmicb.2015.00668.

598 Soong, G., Martin, F.J., Chun, J., Cohen, T.S., Ahn, D.S., Prince, A., 2011. *Staphylococcus*
599 *aureus* protein A mediates invasion across airway epithelial cells through activation of RhoA
600 GTPase signaling and proteolytic activity. *J Biol Chem* 286, 35891–35898. doi:
601 10.1074/jbc.M111.295386.

602 Tajima, A., Seki, K., Shinji, H., Masuda, S., 2006. Inhibition of interleukin-8 production in
603 human endothelial cells by *Staphylococcus aureus* supernatant. *Clin Exp Immunol* 147,
604 148–154. doi: 10.1111/j.1365-2249.2006.03254.x

605 Thammavongsa, V., Kim, H.K., Missiakas, D., Schneewind, O., 2015. Staphylococcal
606 manipulation of host immune responses. *Nat Rev Micro* 13, 529–543. doi:
607 10.1038/nrmicro3521.

608 Tisoncik, J.R., Korth, M.J., Simmons, C.P., Farrar, J., Martin, T.R., Katze, M.G., 2012. Into
609 the eye of the cytokine storm. *Microbiol Mol Biol Rev* 76, 16–32. doi: 10.1128/MMBR.05015-
610 11.

611 Tong, S.Y.C., Davis, J.S., Eichenberger, E., Holland, T.L., Fowler, V.G., 2015.
612 *Staphylococcus aureus* infections: epidemiology, pathophysiology, clinical manifestations,
613 and management. *Clin Microbiol Rev* 28, 603–661. doi: 10.1128/CMR.00134-14.

614 Troeger, C., Forouzanfar, M., Rao, P.C., Khalil, I., Brown, A., Swartz, S., Fullman, N.,
615 Mosser, J., Thompson, R.L., Reiner, R.C., Abajobir, A., Alam, N., Alemayohu, M.A., Amare,
616 A.T., Antonio, C.A., Asayesh, H., Avokpaho, E., Barac, A., Beshir, M.A., Boneya, D.J.,
617 Brauer, M., Dandona, L., Dandona, R., Fitchett, J.R.A., Gebrehiwot, T.T., Hailu, G.B., Hotez,
618 P.J., Kasaeian, A., Khoja, T., Kisseon, N., Knibbs, L., Kumar, G.A., Rai, R.K., Razek, El,
619 H.M.A., Mohammed, M.S.K., Nielson, K., Oren, E., Osman, A., Patton, G., Qorbani, M.,
620 Roba, H.S., Sartorius, B., Savic, M., Shigematsu, M., Sykes, B., Swaminathan, S., Topor-
621 Madry, R., Ukwaja, K., Werdecker, A., Yonemoto, N., Sayed Zaki, El, M., Lim, S.S.,
622 Naghavi, M., Vos, T., Hay, S.I., Murray, C.J.L., Mokdad, A.H., 2017. Estimates of the global,
623 regional, and national morbidity, mortality, and aetiologies of lower respiratory tract
624 infections in 195 countries: a systematic analysis for the Global Burden of Disease Study
625 2015. *Lancet Infect Dis* 17, 1133–1161. doi: 10.1016/S1473-3099(17)30396-1.

626 Urban, T.A., Goldberg, J.B., Forstner, J.F., Sajjan, U.S., 2005. Cable pili and the 22-
627 kilodalton adhesin are required for *Burkholderia cenocepacia* binding to and transmigration
628 across the squamous epithelium. *Infect Immun* 73, 5426–5437. doi: 10.1128/IAI.73.9.5426-
629 5437.2005

630 Urban, T.A., Griffith, A., Torok, A.M., Smolkin, M.E., Burns, J.L., Goldberg, J.B., 2004.
631 Contribution of *Burkholderia cenocepacia* flagella to infectivity and inflammation. *Infect*
632 *Immun* 72, 5126–5134. doi: 10.1128/IAI.72.9.5126-5134.2004.

633 Voyich, J.M., Otto, M., Mathema, B., Braughton, K.R., Whitney, A.R., Welty, D., Long, R.D.,
634 Dorward, D.W., Gardner, D.J., Lina, G., Kreiswirth, B.N., DeLeo, F.R., 2006. Is Pantone-
635 Valentine leukocidin the major virulence determinant in community-associated methicillin-
636 resistant *Staphylococcus aureus* disease? *J Infect Dis* 194, 1761–1770. doi:
637 10.1086/509506

638 Wachsmann, P., Lamprecht, A., 2012. Polymeric nanoparticles for the selective therapy of
639 inflammatory bowel disease. *Methods Enzymol* 508, 377–397. doi: 10.1016/B978-0-12-
640 391860-4.00019-7.

641

642

643

644

645

646

647

648

649 **Table 1 Bacteria strains**

Strain	Characteristics	Reference
MRSA252	Hospital-acquired MRSA, fatal post-op septicemia	Holden et al., 2004
MSSA NCTC 6571	Methicillin-sensitive <i>S. aureus</i>	
MSSA Newman	Isolated from a secondary osteomyelitis infection of a TB patient	Baba et al., 2008
MSSA209		Collins et al., 2010
MRSA41		Collins et al., 2010
MSSA SH1000	Laboratory strain, derivative of NCTC 8325 by complementation of <i>rsbU</i>	O'Neill, 2010
MSSA RN4220	Derivative of NCTC8325-4, with <i>agrA</i> mutation	Nair et al., 2011
MRSA USA400	Community-associated <i>S. aureus</i>	Voyich et al., 2006
<i>B. cenocepacia</i> J2315	Epidemic CF-related ET12 lineage	Holden et al., 2009

650

651

652

653

654

Figure 1 *S. aureus* supernatant has anti-inflammatory on *B. cenocepacia*-induced IL-8 response.

IL-8 production in 16HBE cells treated with *B. cenocepacia* (5 MOI), in the presence or absence of **(A)** 10% *S. aureus* supernatants, or **(B)** heat-killed *S. aureus* (5 MOI) for 24 hours. The experiments were validated using **(C)** 10% supernatants from *S. aureus* isolates or **(D)** heat-killed *S. aureus* and *B. cenocepacia* (MOI 5) co-stimulation for 24 hours. In **(A)** and **(C)**, 10% LB was used as a control. Data represent mean \pm SEM of duplicate samples from three identical experiments. Significance was compared to *B. cenocepacia*-treated cells using 2-way ANOVA followed by Dunnett's multiple comparison test (*, $p < 0.05$).

(E) Cytotoxicity of *S. aureus* supernatants on 16HBE cells was assessed using MTT assay. The control was treated with 10% LB. Results are presented as a percent of the control, mean \pm SEM from three independent experiments. Significance was compared to the control using 1-way ANOVA followed by Dunnett's multiple comparison test.

Figure 2 Effect of *S. aureus* supernatants on by *B. cenocepacia*-activated MAPK and NF- κ B signalling.

16HBE cells were infected by *B. cenocepacia* unwashed culture at an MOI of 5 for the indicated time. TNF α (20 ng/mL) for 5 minutes was used as the positive control. **(A)** Western blot analysis for levels of phosphorylated p38, phosphorylated Erk, phosphorylated p65, and p38 in cell lysate followed by *B. cenocepacia* stimulation. Phosphorylation of p38 **(B)**, ERK **(C)**, p65 **(D)** of cell lysates were measured by western blot and normalised against total p38.

16HBE cells were treated with 10% LB as the control, *B. cenocepacia* (5 MOI), with the presence or absence of 10% *S. aureus* supernatant for 2 hours. **(E)** Western blot analysis for levels of phosphorylated p38, phosphorylated Erk, phosphorylated p65, and p38 in cell lysate following *S. aureus* supernatants and *B. cenocepacia* co-stimulation. Phosphorylation of p38

(F), Erk (G) and p65 (H) of cell lysates were measured using western blot and normalised against total p38.

Protein bands representative of three independent experiments. Data are shown as mean \pm SEM. *, $p < 0.05$ significant reduction compared to *B. cenocepacia*-treated group using 2-way ANOVA followed by a one-tailed Dunnett's multiple comparisons test.

Figure 3 Characterisation of anti-inflammatory components in *S. aureus* supernatants.

16HBE cells were treated with *B. cenocepacia* (5 MOI), in the presence or absence of 10% (A) heat-treated supernatants, (B) proteinase K (PK)-treated *S. aureus* supernatants, (C) *S. aureus* supernatant fractions containing metabolites of >10 kDa, >3 kDa, and <3 kDa for 24 hours. IL-8 production was analysed using ELISA. Data represent mean \pm SEM from 3 independent experiments performed in duplicate. *, $p < 0.05$ significance compared to *B. cenocepacia*-treated group using 2-way ANOVA followed by Dunnett's multiple comparisons test.

(D) 16HBE cells were co-treated with IL-1 (1 ng/mL) / TNF α (10 ng/mL) and 10% *S. aureus* supernatants for 24 hours. *B. cenocepacia* (5 MOI) was the positive control. Data represent mean \pm SEM from 3 independent experiments performed in duplicate. *, $p < 0.05$ compared to IL-1 β / TNF α -treated group or with IL-1/ TNF α -treated group using 2-way ANOVA followed by Tukey's multiple comparisons test.




Figure 1




Figure 2




Figure 3